

# An International Journal of Research in AYUSH and Allied Systems

Case Study

# AYURVEDIC MANAGEMENT OF RAKTAPRADARA (HEAVY MENSTRUAL BLEEDING)

## Pratibha Bahogona

Assistant Professor, Dept. of Sharir kriya vigyan, Dronacharya Ayurvedic Educational Institute & Hospital, Saharanpur, Uttar Pradesh India.

#### Article info

Article History: Received: 19-01-2025 Accepted: 27-02-2025 Published: 20-03-2025

#### **KEYWORDS:**

Raktapradara, Menorrhagia, Dysfunctional uterine bleeding, Shamana therapy, Shodhana therapy. ABSTRACT

*Raktapradara*, as referenced in the Charak Samhita, signifies excessive menstrual bleeding resulting from the imbalances of the Rakta and Pitta doshas. This condition is akin to menorrhagia or dysfunctional uterine bleeding recognized in contemporary medicine. Various dietary and lifestyle factors can exacerbate the issue, leading to heightened blood loss, fatigue, and irregular menstrual cycles. Ayurvedic approaches advocate a comprehensive treatment plan that includes herbal remedies, *Panchakarma* therapy, dietary changes, and lifestyle modifications aimed at restoring menstrual health. The objective of this study is to assess the effectiveness of Ayurvedic treatment in addressing Raktapradara through a detailed case analysis, focusing on the role of herbal interventions and Panchakarma therapy in normalizing menstrual cycles and enhancing overall health. In the methods section, a 32-year-old woman experiencing heavy menstrual bleeding for the duration of three months received Ayurvedic treatment. This included a regimen of oral medications, such as Chandraprava Vati, Sonitargala Rasa, Nagakesara Churna, and Ashokarishta, along with Panchakarma therapy involving Virechana with Haritaki Churna, alongside specific dietary and lifestyle modifications. The treatment lasted for three months, with evaluations based on factors including menstrual flow, cycle consistency, hemoglobin levels, and general health improvements. The results indicated that the patient experienced notable enhancements after the three-month treatment period.

#### **INTRODUCTION**

In the Charak Samhita, "*Raktapradar*" signifies the excessive discharge of menstrual blood caused by the *Pradirna* of *Raja*; so, it is termed *Pradar*. *Raktapradar* is more strongly linked to heavy menstrual bleeding in terms of its pathogenesis. Normal menstrual blood loss varies between 50 and 80 ml and should not exceed 100ml. Menorrhagia refers to excessive vaginal bleeding, defined by an increased volume, extended duration, or both. "Dysfunctional uterine bleeding" denotes abnormal menstrual bleeding that cannot be ascribed to any discernible structural or systemic disorder. Utilized specifically when menorrhagia is not associated with any

#### Access this article online



abnormalities of the genital tract or systemic or endocrine problems. According to the Charak Samhita, a woman who excessively ingests *Lavan, Amla, Guru, Katu, Vidāhi, Snigdha, Mamsa, Krśara, Pāyasa,* curd, *Sukta, Mastu,* and *Madya* suffers from the vitiation of *Rakta,* leading to an increase in its volume; thus, the elevation of *Raja* is due to its integration with augmented blood. This condition is commonly known as *Pradara* due to the profuse hemorrhage associated with it. The therapeutic approach for this ailment should incorporate *Raktasthapak* alongside *Vata Pitta Shamak.* 

#### **AIMS AND OBJECTIVES**

- To evaluate the effectiveness of Ayurvedic treatment in managing *Raktapradara*.
- To assess the role of herbal and *Panchakarma* interventions in restoring menstrual health.
- To provide a safe and sustainable alternative to conventional treatments.

## AYUSHDHARA, 2025;12(1):127-130

#### MATERIALS AND METHODS

Study Design: Single case study

Patient Selection: A 32-year-old female with excessive and prolonged menstrual bleeding.

Intervention: Ayurvedic treatment including oral medications, dietary modifications, and Panchakarma therapy.

**Duration: 3 months** 

Assessment Criteria: Reduction in bleeding duration and quantity, improvement in general health, and regularization of menstrual cycles.

## **Case Presentation**

A 32-year-old married female came to hospital OPD & presented with complaints of excessive menstrual bleeding for 3 months, associated with weakness, dizziness, and mild lower abdominal pain. The patient had no history of systemic diseases, but stress and dietary irregularities were noted.

N/H/O- HTN, DM, Thyroid etc.

### **Personal History**

Diet: Mixed

Appetite: Good

### **Treatment Protocol**

Intern

Sleep: Disturbed **Bowel Habits: Irregular** Micturition: 6-7/day **Menstrual History** Menarche - 13 years Interval: 22-24 days Duration of Bleeding: 10-12 days Amount: 4-5 pads/ day Flow: Heavy, associated with clots. Pain: Mild lower abdominal discomfort. **Clinical Examination** Pallor: Present Pulse: 70/min. BP: 120/70 mmHg Abdomen: Soft. non-tender **Obstetric history:** G1P1L1A0 (FTNVD) Investigation: Hb: 8gm USG Pelvis & Abd- No Finding Urine: NAD

al Intervention			
S.No	Drugs	Dose	Duration
1	Chandraprava Vati	2 BD	3 months
2	Sonitargala Rasa	1 BD	3 months
3	Nagakesara Churna	5gm, BD	Bleeding time, 10 days each cycle
4	Ashokarista	20ml BD with equal water	3 months

# Panchakarma Therapy

Virechana with Haritaki churna 5gm with lukewarm water 5 days.

# **Dietary Modifications**

Inclusion of cooling and *Pitta*-pacifying foods like milk, ghee, and fresh fruits.

Avoidance of spicy, sour, and heat-inducing foods.

## **Lifestyle Advice**

Stress management through yoga and meditation. Regular physical activity like walking.

### **OBSERVATION AND RESULTS**

After three months of treatment, the patient showed significant improvement:

Menstrual Flow: Reduced to 4-5 days with moderate bleeding.

Cycle Regularity: Improved to 28 days.

General Health: Increased energy levels, reduced dizziness and pallor.

No Adverse Effects: The treatment was well tolerated.

# **Statistical Data**

- Pre-treatment average blood loss: 250-300ml per cvcle
- Post-treatment average blood loss: 100-150ml per cvcle
- Hemoglobin levels before treatment: 9.2 g/dL
- Hemoglobin levels after treatment: 11.5 g/dL
- Reduction in fatigue scores: From 8/10 to 3/10 on a patient-reported scale.



#### DISCUSSION

The Ayurvedic approach to managing Raktapradara, or heavy menstrual bleeding, emphasizes restoring balance to the aggravated Pitta and *Rakta doshas*, which are the primary contributors to increased blood flow and associated symptoms. As outlined in the Charak Samhita, an imbalance in these Doshas can result in heightened menstrual bleeding, often accompanied by heat, inflammation, and irregular cycles. The treatment methodology in Avurveda includes the incorporation of *Pitta-shamak* (*Pitta*-pacifying) herbs, hemostatic agents (*Raktasthambhak*), uterine tonics, and detoxification therapies, all aimed at regulating the menstrual cycle and achieving systemic equilibrium. Among the various remedies. Ashokarishta is particularly significant in the treatment of Raktapradara. This herbal formulation, made from the bark of Saraca asoca, is known for its astringent, hemostatic, and uterine tonic effects. It enhances uterine muscle strength, regulates menstrual flow, and supports hormone balance, which is vital for women's reproductive health. The formulation is enriched with bioactive compounds such as flavonoids and tannins that help reduce excessive bleeding and improve endometrial integrity. In addition, Sonitargala Rasa, a herbo-mineral preparation, works in synergy with Ashokarishta to further promote hemostasis and minimize blood loss. A fundamental component of this treatment protocol is Virechana, a detoxification therapy that utilizes Haritaki Churna. Virechana acts as a purging treatment that eliminates excess *Pitta* from the body, alleviating systemic inflammation and correcting hormonal discrepancies. Haritaki (Terminalia chebula) is recognized for its gentle laxative and detoxifying properties, which enhance digestion and metabolism- key factors in regulating menstrual health. By clearing excess heat and toxins, Virechana aids in restoring homeostasis and bolstering

overall vitality. In addition to these therapeutic methods, dietary and lifestyle changes play a crucial role in the Ayurvedic management of *Raktapradara*. Patients are encouraged to consume cooling and nourishing foods, such as milk, ghee, fresh fruits, and fiber-rich vegetables, which help balance *Pitta* and replenish essential nutrients. In contrast, the intake of spicy, sour, and heat-inducing foods is limited, as these can exacerbate *Pitta* and worsen menstrual symptoms. Furthermore, practices like yoga and meditation are recommended to alleviate stress, which is integral to maintaining endocrine and reproductive health.

The clinical enhancements noted in the patient. such as a decrease in menstrual bleeding, elevated hemoglobin levels, and increased energy, illustrate the success of this holistic Ayurvedic method. The lack of negative side effects, along with the overall improvement in menstrual health, underscores the promise of Avurveda as a safe and sustainable option compared to traditional hormonal or surgical interventions heavv menstrual for bleeding. Nonetheless, it is crucial to conduct further large-scale clinical trials to confirm these results and to develop standardized treatment guidelines for the Ayurvedic management of Raktapradara.

#### CONCLUSION

This study illustrates the efficacy of Ayurvedic Raktapradara treatment in addressing fheavv menstrual bleeding) by employing a comprehensive combines herbal approach that remedies. Panchakarma therapies, dietary changes, and lifestyle adjustments. The findings indicate that utilizing a mix of these strategies can considerably enhance menstrual health by tackling the fundamental causes of excessive bleeding, re-establishing hormonal balance, and improving overall wellness. The incorporation of *Pitta*-pacifying and hemostatic herbs, including

#### AYUSHDHARA, 2025;12(1):127-130

Ashokarishta and Sonitargala Rasa, was essential in managing excessive menstrual blood flow, while Virechana therapy utilizing Haritaki Churna aided in detoxifying the body and correcting *Doshic* imbalances. In addition to mitigating blood loss, the treatment approach resulted in better hemoglobin levels, heightened energy, and diminished fatigue, reflecting a positive influence on systemic health. The patient reported improved regularity of menstrual cycles, indicating that Ayurvedic treatment not only alleviates symptoms but also fosters long-term menstrual health by addressing foundational Doshic imbalances. The lack of negative side effects in this case further confirms the safety and effectiveness of Ayurvedic methods as a practical alternative to conventional hormonal or surgical interventions, which frequently have considerable side effects. Additionally, this study highlights the significance of dietary and lifestyle adjustments in treating menstrual disorders. By integrating cooling and nutrient-dense foods, steering clear of spicy and heat-generating diets, and implementing stress-reduction strategies such as voga and meditation, the patient noted enhanced physical and mental health. These aspects play a role in sustaining reproductive health and decreasing the frequency of menstrual irregularities. Although this individual case study presents promising outcomes, it accentuates the necessity for larger clinical trials to corroborate these results and formulate standardized Ayurvedic treatment protocols for *Raktapradara*. Future investigations should prioritize comparative analyses between Ayurvedic and allopathic treatments, long-term assessments, and the examination of the mechanisms behind herbal formulas in managing menstrual issues. The incorporation of Ayurveda into standard healthcare practices has the potential to offer a safe, natural, and sustainable solution for women's health challenges, thereby fostering holistic wellness and enhancing quality of life.

## REFERENCES

- 1. Edited by- Vaidya shastri shree laxmipati Yogratnakar: vidyotini Hindi Tikta Samhita Chaukhambha surbharti prakashan, Varanasi. Year-2015 Edition.
- 2. Edited by- Tripathi Bramhanand Charaksamhita Chaukhambha Bharati academy, Varanasi. Year-2013 Edition.
- 3. Edited by- Dr. Sharma Anantram sushrut samhita Volume-2 Chaukhamba surbharati prakashan, Varanasi. Year-2009 Edition.
- 4. Edited by- Tiwari Premvati Ayurvediya Prasutitantra evam Streeroga part 2, Chaukhamaba orientalia, Varanasi. Year- 2014 Edition.
- 5. Edited by- Ayurvedacharya Sharma Pandit heeralal Sharangdhar Samhita Chaukhamba orientalia, Varanasi Year-2008 Edition.
- 6. D C Dutta, Text Book of Gynecology, 6<sup>th</sup> Edition, Jaypee Brothers Medical Publisher (P) Limited, Chapter 14, p185 3.

Cite this article as: Pratibha Bahogona. Ayurvedic Management of Raktapradara (Heavy Menstrual Bleeding). AYUSHDHARA, 2025;12(1):127-130. https://doi.org/10.47070/ayushdhara.v12i1.1946 Source of support: Nil, Conflict of interest: None Declared \*Address for correspondence Dr. Pratibha Bahogona Assistant Professor, Dept. of Sharir Kriya Vigyan, Dronacharya Ayurvedic Educational Institute & Hospital, Saharanpur, Uttar Pradesh, India. Email: Kalpanabahogona@gmail.com

Disclaimer: AYUSHDHARA is solely owned by Mahadev Publications - A non-profit publications, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. AYUSHDHARA cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of AYUSHDHARA editor or editorial board members.